

In vivo Animal Models

## Huntington's Disease



## BACHD Transgenic Rat Model

BACHD rats overexpress full-length human mHTT with 97 alternating CAA/CAG repeats on a bacterial artificial chromosome (BAC) that are sufficient to produce the Huntington's disease (HD) phenotype:

HTT aggregates

## polyQ HTT

Figure 1:

1000.000 1000.0000 1000.000 1000.00000 1000.0000 1000.0000 1000000 10000000 1000000

Figure 1: Quantification of polyQ HTT by sandwich immunosorbent assay using the MesoScale Discovery platform. Whole brain lysate of a 6 months old hemizygous rat and a non-transgenic littermate were used in a dilution series of 1:10, 1: 40; 1:160; 1:640 and 1:2560.

Yu-Taeger L, Petrasch-Parwez E, Osmand AP, Redensek A, Metzger S, Clemens LE, Park L, Howland D, Calaminus C, Gu X Pichler B, Yang XW, Riess O, Nguyen HP. A novel BACHD transgenic rat exhibits characteristic neuropathological features of Huntington disease. J Neurosci. 2012 Oct 31;32(44):15426-38.

Scantox Discovery

## Scantox Group, HQ

Hestehavevej 36A, Ejby DK – 4623 Lille Skensved clientservice@scantox.com www.scantox.com +45 5686 1500 © Scantox A/S Scantox is a registered trademark of Scantox A/S.



126-36-HuntingtonsBACHD-EN (04/2024)