



## BTBR T+ tf/J Inbred Mouse Model


BTBR T+ tf/J mice are an inbred strain completely missing the corpus callosum and a severely reduced hippocampal commissure. As negative control, wild type inbred mice (e. g. C57BL/6) can be used.

- Social behavior and interaction deficits
- Repetitive behavior
- Reduced anxiety
- Spatial learning deficits
- Nest building and muscle strength deficits


**Figure 1:** Three chamber social interaction test of BTBR mice. Time spent sniffing in seconds at the unfamiliar mouse or the empty cup during social approach (A) and the time spent sniffing in seconds at the familiar or the unfamiliar mouse during social recognition (B). n = 14 - 15; males only; mean + SEM. Two-way ANOVA with Bonferroni's post hoc test. \*p<0.05; \*\*p<0.01.

Figure 1

### Social Approach Time Sniffing




### Recognition Time Sniffing




**Figure 2:** Autogrooming behavior in BTBR mice (A). Mean duration of grooming bouts. Time spent in the open arms of the elevated plus maze test (B). n = 14 - 15; males only; mean + SEM; unpaired t-test; \*\*p<0.01; \*\*\*p<0.001.

Figure 2

### Grooming Bouts



### Time in Open Arms



e. g.: McFarlane et al. Autism-like behavioral phenotypes in BTBR T+tf/J mice. *Genes Brain Behav.* 2008 Mar;7(2):152-63.

